Neuroscience Letters, 96 (1989) 356–359
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, K., see Ikeda, M., 202
Adams, S.A., see Saper, CB., 29
Agid, Y., see Hirsch, E.C., 145
Ahmed, Z., see Lewis, C.A., 185
Allen, G.S., see Whetsell Jr., W.O., 18
Altar, C.A. and Marien, M.R., Preservation of dopamine release in the denervated striatum, 329
Alving, K., see Martling, C.-R., 306
Ates, N., see Ziylan, Y.Z., 179

Benabid, A.L. and Jeaugey, L., Cells of the rat lateral habenula respond to high-threshold somatosensory inputs, 289
Benoliel, J.J., see Pohl, M., 102
Bocchieri, P., see Marchi, M., 229
Bonnet, J.-J., see Thibaut, F., 335
Bourgoin, S., see Pohl, M., 102
Bourre, J.M., see Souyri, F., 351
Braak, H., see Ohm, T.G., 60
Butcher, L.L., see Woolf, N.J., 277
Bähr, M., Vanselow, J. and Thanos, S., Ability of adult rat ganglion cells to regrow axons in vitro can be influenced by fibroblast growth factor and gangliosides, 197

Cesselin, F., see Pohl, M., 102
Chahl, L.A., Kavanagh, C.A. and Corlis, A., Enhancement by haloperidol of the locomotor response induced by naloxone in morphine-treated guinea pigs, 213
Chan, S.H.H., see Lui, P.-W., 114
Chen, H., Singh, Y.N. and Dryden, W.F., Transduction mechanism involving the presynaptic adenosine receptor at mouse motor nerve terminals, 318
Chesselet, M.-F. and Robbins, E., Regional differences in substance P-like immunoreactivity in the striatum correlate with levels of pre-protachykinin mRNA, 47
Coggeshall, R.E., see Westlund, K.N., 13
Corlis, A.,see Chahl, L.A., 213
Corvetti, G., see Levi, A.C., 271
Costa, E., see Danysz, W., 340
Costenin, J., see Thibaut, F., 335

D'Angio, M. and Scatton, B., Feeding or exposure to food odors increases extracellular DOPAC levels (as measured by in vivo voltammetry) in the prefrontal cortex of food-deprived rats, 223
Dahl, D., see Gilad, G.M., 7
Danysz, W., Fadda, E., Wroblewski, J.T. and Costa, E., Kynurenate and 2-amino-5-phosphonovalerate interact with multiple binding sites of the *N*-methyl-D-aspartate-sensitive glutamate receptor domain, 340
DeMattei, M., see Levi, A.C., 271
Deschenes, N.L., see Ransom, R.W., 323
Diaz, J., Molgó, J. and Pécot-Dechavassine, M., Sprouting of frog motor nerve terminals after long-term paralysis by botulinum type A toxin, 127
Dietzel, I.D., see Gottmann, K., 173
Doble, A., see Hubert, J.P., 345
Doherty, P. and Walsh, F.S., K-252a specifically inhibits the survival and morphological differentiation of NGF-dependent neurons in primary cultures of human dorsal root ganglia, 1
Dryden, W.F., see Chen, H., 318
Dubner, R., see Nahin, R.L., 247
Dubuc, R., see Fritzsch, B., 241
Dumas, J., see Parent, A., 140
Duyckaerts, C., see Hirsch, E.C., 145

Eisenberg, B. and Messer, A., Toxic/clonic seizures in a mouse mutant carrying the weaver gene, 168
Ewert, J.-P., see Gonzalez-Lima, F., 151

Faber, D.S., see Lewis, C.A., 185
Fadda, E., see Danysz, W., 340
Fariello, R.G., see Levi, A.C., 271
Felix, D., see Imboden, H., 42
Fetz, E.E., see Smith, W.S., 76
Filion, M., see Parent, A., 140
Finger, T.E., see Mahalik, T.J., 253
Finkenstädt, T., see Gonzalez-Lima, F., 151
Fritzsch, B., Dubuc, R., Ohta, Y. and Grillner, S., Efferents to the labyrinth of the river lamprey (*Lampetra fluviatilis*) as revealed with retrograde tracing techniques, 241

Gamble, G.D. and Milne, R.J., Repeated exposure to sham testing procedures reduces reflex withdrawal and hot-plate latencies: attenuation of tonic descending inhibition?, 312

Neuroscience Letters, 96 (1989) 356–359
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, K., see Ikeda, M., 202
Adams, S.A., see Saper, CB., 29
Agid, Y., see Hirsch, E.C., 145
Ahmed, Z., see Lewis, C.A., 185
Allen, G.S., see Whetsell Jr., W.O., 18
Altar, C.A. and Marien, M.R., Preservation of dopamine release in the denervated striatum, 329
Alving, K., see Martling, C.-R., 306
Ates, N., see Ziylan, Y.Z., 179

Benabid, A.L. and Jeaugey, L., Cells of the rat lateral habenula respond to high-threshold somatosensory inputs, 289
Benoliel, J.J., see Pohl, M., 102
Bocchieri, P., see Marchi, M., 229
Bonnet, J.-J., see Thibaut, F., 335
Bourgoin, S., see Pohl, M., 102
Bourre, J.M., see Souyri, F., 351
Braak, H., see Ohm, T.G., 60
Butcher, L.L., see Woolf, N.J., 277
Bähr, M., Vanselow, J. and Thanos, S., Ability of adult rat ganglion cells to regrow axons in vitro can be influenced by fibroblast growth factor and gangliosides, 197

Cesselin, F., see Pohl, M., 102
Chahl, L.A., Kavanagh, C.A. and Corlis, A., Enhancement by haloperidol of the locomotor response induced by naloxone in morphine-treated guinea pigs, 213
Chan, S.H.H., see Lui, P.-W., 114
Chen, H., Singh, Y.N. and Dryden, W.F., Transduction mechanism involving the presynaptic adenosine receptor at mouse motor nerve terminals, 318
Chesselet, M.-F. and Robbins, E., Regional differences in substance P-like immunoreactivity in the striatum correlate with levels of pre-protachykinin mRNA, 47
Coggeshall, R.E., see Westlund, K.N., 13
Corlis, A.,see Chahl, L.A., 213
Corvetti, G., see Levi, A.C., 271
Costa, E., see Danysz, W., 340
Costenin, J., see Thibaut, F., 335

D'Angio, M. and Scatton, B., Feeding or exposure to food odors increases extracellular DOPAC levels (as measured by in vivo voltammetry) in the prefrontal cortex of food-deprived rats, 223
Dahl, D., see Gilad, G.M., 7
Danysz, W., Fadda, E., Wroblewski, J.T. and Costa, E., Kynurenate and 2-amino-5-phosphonovalerate interact with multiple binding sites of the *N*-methyl-D-aspartate-sensitive glutamate receptor domain, 340
DeMattei, M., see Levi, A.C., 271
Deschenes, N.L., see Ransom, R.W., 323
Diaz, J., Molgó, J. and Pécot-Dechavassine, M., Sprouting of frog motor nerve terminals after long-term paralysis by botulinum type A toxin, 127
Dietzel, I.D., see Gottmann, K., 173
Doble, A., see Hubert, J.P., 345
Doherty, P. and Walsh, F.S., K-252a specifically inhibits the survival and morphological differentiation of NGF-dependent neurons in primary cultures of human dorsal root ganglia, 1
Dryden, W.F., see Chen, H., 318
Dubner, R., see Nahin, R.L., 247
Dubuc, R., see Fritzsch, B., 241
Dumas, J., see Parent, A., 140
Duyckaerts, C., see Hirsch, E.C., 145

Eisenberg, B. and Messer, A., Toxic/clonic seizures in a mouse mutant carrying the weaver gene, 168
Ewert, J.-P., see Gonzalez-Lima, F., 151

Faber, D.S., see Lewis, C.A., 185
Fadda, E., see Danysz, W., 340
Fariello, R.G., see Levi, A.C., 271
Felix, D., see Imboden, H., 42
Fetz, E.E., see Smith, W.S., 76
Filion, M., see Parent, A., 140
Finger, T.E., see Mahalik, T.J., 253
Finkenstädt, T., see Gonzalez-Lima, F., 151
Fritzsch, B., Dubuc, R., Ohta, Y. and Grillner, S., Efferents to the labyrinth of the river lamprey (*Lampetra fluviatilis*) as revealed with retrograde tracing techniques, 241

Gamble, G.D. and Milne, R.J., Repeated exposure to sham testing procedures reduces reflex withdrawal and hot-plate latencies: attenuation of tonic descending inhibition?, 312

Garbarino, L., see Marchi, M., 229
Gilad, G.M., Gilad, V.H. and Dahl, D., Expression of neurofilament immunoreactivity in developing rat cerebellum in vitro and in vivo, 7
Gilad, V.H., see Gilad, G.M., 7
Golden, G.T., see Levi, A.C., 271
Gonzalez-Lima, F., Finkenstädt, T. and Ewert, J.-P., Neural substrates for long-term habituation of the acoustic startle reflex in rats: a 2-deoxyglucose study, 151
Gottmann, K., Dietzel, I.D. and Lux, H.D., Proton-induced chloride current and voltage-activated Na^+ and the Ca^+ current in embryonic neurons from the medicinal leech (*Hirudo medicinalis*), 173
Grace, A.A., see Hollerman, J.R., 82
Graybiel, A.M., see Hirsch, E.C., 145
Grillner, S., see Fritzsch, B., 241

Hamon, M., see Pohl, M., 102
Harding, J.W., see Imboden, H., 42
Hersh, L.B., see Hirsch, E.C., 145
Hervonen, A., see Sirviö, J., 218
Hirsch, E.C., Graybiel, A.M., Hersh, L.B., Duyckaerts, C. and Agid, Y., Striosomes and extrastriosomal matrix contain different amounts of immunoreactive choline acetyltransferase in the human striatum, 145
Hollerman, J.R. and Grace, A.A., Acute haloperidol administration induces depolarization block of nigral dopamine neurons in rat after partial dopamine lesions, 82
Holmes, H.R., see Saper, C.B., 29
Hubert, J.P. and Doble, A., Ibotenic acid stimulates D-[^{3}H]aspartate release from cultured cerebellar granule cells, 345
Hurley, K.M., see Saper, C.B., 29
Hylden, J.L.K., see Nahin, R.L., 247

Iadarola, M.J., see Nahin, R.L., 247
Ikeda, M., Nakazawa, T., Abe, K., Kaneko, T. and Yamatsu, K., Extracellular accumulation of glutamate in the hippocampus induced by ischemia is not calcium dependent – in vitro and in vivo evidence, 202
Ikeda, S.R. and Schofield, G.G., Somatostatin cyclic octapeptide analogs which preferentially bind to SOMa receptors block a calcium current in rat superior cervical ganglion neurons, 283
Ikuta, F., see Wakabayashi, K., 259
Imboden, H., Harding, J.W. and Felix, D., Hypothalamic angiotensinergic fibre systems terminate in the neurohypophysis, 42
Inagaki, M., Takeshita, K., Nakao, S., Shiraishi, Y. and Oikawa, T., An electrophysiologically defined trigemino-reticulo-facial pathway related to the blink reflex in the cat, 64

Jacobs, R.W., see Woolf, N.J., 277
Jeaugey, L., see Benabid, A.L., 289
Jeftinija, S., Excitatory transmission in the dorsal horn is a part mediated through APV-sensitive NMDA receptors, 191

Kakei, S., see Shinoda, Y., 163
Kaneko, K., see Ikeda, M., 202
Kani, K., see Sakai, M., 157
Kavanagh, C.A., see Chahl, L.A. 213
Kemali, D., see Kemali, M., 235
Kemali, M., Monteleone, P., Maj, M., Milici, N. and Kemali, D., Lithium decreases retinal melatonin levels in the frog, 235
Kunkel, D.D., see Scharfman, H.E., 23

Lacroix, J.S., see Martling, C.-R., 306
Leahy, K.M., see Saper, C.B., 29
Lee, T.-Y., see Lui, P.-W., 114
Levi, A.C., DeMattei, M., Ravazzani, R., Corvetti, G., Golden, G.T. and Fariello, R.G., Effects of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) on ultrastructure of nigral neuromelanin in *Macaca fascicularis*, 271
Lewis, C.A., Ahmed, Z. and Faber, D.S., Characteristics of glycine-activated conductances in cultured medullary neurons from embryonic rat, 185
Lidsky, T.I., see Manetto, C., 295
Lui, P.-W., Lee, T.-Y. and Chan, S.H.H., Involvement of locus coeruleus and noradrenergic neurotransmission in fentanyl-induced muscular rigidity in the rat, 114
Lukas, R.J., see Oswald, R.E., 207
Lundberg, J.M., see Martling, C.-R., 306
Lux, H.D., see Gottmann, K., 173

Macdonald, J.F., see Mody, I., 70
Macdonald, R.L., see Twyman, R.E., 89
Magnusson, K. and Wieloch, T., Impairment of protein ubiquitination may cause delayed neuronal death, 264
Mahalik, T.J., Stromberg, I., Finger, T.E. and Olson, L., Abnormal expression of tyrosine hydroxylase-like immunoreactivity in intraocular transplants of rat caudate nucleus, 253
Maj, M., see Kemali, M., 235

Manetto, C. and Lidsky, T.I., The effects of movements on caudate sensory responses, 295
Marchi, M., Bocchieri, P., Garbarino, L. and Raiteri, M., Muscarinic inhibition of endogenous glutamate release from rat hippocampus synaptosomes, 229
Marien, M.R., see Altar, C.A., 329
Marson, L., Evidence for colocalization of substance P and 5-hydroxytryptamine in spinally projecting neurons from the rat medulla oblongata, 54
Martling, C.-R., Matran, R., Alving, K., Lacroix, J.S. and Lundberg, J.M., Vagal vasodilatory mechanisms in the pig bronchial circulation preferentially involves sensory nerves, 306
Martínez-Murillo, R., Villalba, R.M. and Rodrigo, J., Electron microscopic localization of cholinergic terminals in the rat substantia nigra: an immunocytochemical study, 121
Matran, R., see Martling, C.-R., 306
Matthies, H., see Reymann, K.G., 96
Matthies, H.K., see Reymann, K.G., 96
Mauborgne, A., see Pohl, M., 102
McNeill, D.L., see Westlund, K.N., 13
Messer, A., see Eisenberg, B., 168
Milici, N., see Kemali, M., 235
Milne, R.J., see Gamble, G.D., 312
Miyamoto, T., see Okada, Y., 108
Mody, I., Salter, M.W. and MacDonald, J.F., Whole-cell voltage-clamp recordings in granule cells acutely isolated from hippocampal slices of adult or aged rats, 70
Moga, M.M., see Saper, C.B., 29
Molgó, J., see Diaz, J., 127
Monteleone, P., see Kemali, M., 235

Nagatsu, I., see Sakai, M., 157
Nahin, R.L., Hylden, J.L.K., Iadarola, M.J. and Dubner, R., Peripheral inflammation is associated with increased dynorphin immunoreactivity in both projection and local circuit neurons in the superficial dorsal horn of the rat lumbar spinal cord, 247
Nakao, S., see Inagaki, M., 64
Nakazawa, T., see Ikeda, M., 202
Narici, L., see Rossini, P.M., 300
Naus, C.C.G., Development of somatostatin-like immunoreactivity in the hippocampal formation of normal and reeler mice, 133
Needleman, P., see Saper, C.B., 29

Ohama, E., see Wakabayashi, K., 259
Ohm, T.G. and Braak, H., Auditory brainstem nuclei in Alzheimer's disease, 60
Ohta, Y., see Fritzsch, B., 241
Oikawa, T., see Inagaki, M., 64
Okada, Y. and Miyamoto, T., Formation of long-term potentiation in superior colliculus slices from the guinea pig, 108
Olson, L., see Mahalik, T.J., 253
Oswald, R.E., Papke, R.L. and Lukas, R.J., Characterization of nicotinic acetylcholine receptor channels of the TE671 human medulloblastoma clonal line, 207

Papke, R.L., see Oswald, R.E., 207
Parent, A., Smith, Y., Filion, M. and Dumas, J., Distinct afferents to internal and external pallidal segments in the squirrel monkey, 140
Peresson, M., see Rossini, P.M., 300
Pohl, M., Mauborgne, A., Bourgoin, S., Benoliel, J.J., Hamon, M. and Cesselin, F., Neonatal capsaicin treatment abolishes the modulations by opioids of substance P release from rat spinal cord slices, 102
Potter, W.Z., see Saito, K., 35
Pécot-Dechavassine, M., see Diaz, J., 127

Raiteri, M., see Marchi, M., 229
Ransom, R.W. and Deschenes, N.L., Glycine modulation of NMDA-evoked release of [^{3}H]acetylcholine and [^{3}H]dopamine from rat striatal slices, 323
Ravazzani, R., see Levi, A.C., 271
Reymann, K.G., Matthies, H.K., Schulzeck, K. and Matthies, H., *N*-Methyl-D-aspartate receptor activation is required for the induction of both early and late phases of long-term potentiation in rat hippocampal slices, 96
Riekkinen, P.J., see Sirviö, J., 218
Robbins, E., see Chesselet, M.-F., 47
Rodrigo, J., see Martínez-Murillo, R., 121
Rogers, C.J., see Twyman, R.E., 89
Romani, G.L., see Rossini, P.M., 300
Rossini, P.M., Narici, L., Romani, G.L., Peresson, M., Torrioli, G. and Traversa, R., Simultaneous motor output and sensory input: cortical interference site resolved in humans via neuromagnetic measurements, 300

Saavedra, J.M., see Saito, K., 35
Saito, K., Torda, T., Potter, W.Z. and Saavedra, J.M., Characterization of β_1- and β_2-adrenoreceptor subtypes in the rat sinoatrial node and

stellate ganglia by quantitative autoradiography, 35
Sakai, M., Kani, K., Yoshida, M. and Nagatsu, I., Dopaminergic cells in the superior cervical ganglion of the rat: light and electron microscopic study using an antibody against dopamine, 157
Salter, M.W., see Mody, I., 70
Saper, C.B., Hurley, K.M., Moga, M.M., Holmes, H.R., Adams, S.A., Leahy, K.M. and Needleman, P., Brain natriuretic peptides: differential localization of a new family of neuropeptides, 29
Scatton, B., see D'Angio, M., 223
Scharfman, H.E., Kunkel, D.D. and Schwartzkroin, P.A., Intracellular dyes mask immunoreactivity of hippocampal interneurons, 23
Schofield, G.G., see Ikeda, S.R., 283
Schulzeck, K., see Reymann, K.G., 96
Schwartzkroin, P.A., see Scharfman, H.E., 23
Shinoda, Y. and Kakei, S., Distribution of terminals of thalamocortical fibers originating from the ventrolateral nucleus of the rat thalamus, 163
Shiraishi, Y., see Inagaki, M., 64
Singh, Y.N., see Chen, H., 318
Sirviö, J., Riekkinen, P.J. and Hervonen, A., Age-dependence of the solubility fractions of acetylcholinesterase in the cerebral cortex and cerebellum of the rat, 218
Smith, W.S. and Fetz, E.E., Effects of synchrony between primate corticomotoneuronal cells on post-spike facilitation of muscles and motor units, 76
Smith, Y., see Parent, A., 140
Souyri, F. and Bourre, J.M., Altered metabolism of rat contralateral sciatic nerve after microinjection into the endoneurium of the ipsilateral sciatic nerve, 351
Stromberg, I., see Mahalik, T.J., 253

Takahashi, H., see Wakabayashi, K., 259
Takeshita, K., see Inagaki, M., 64
Thanos, S., see Bähr, M., 197
Thibaut, F., Bonnet, J.-J. and Costentin, J., Time coure of postmortem modifications in synaptosomal [^{3}H]dopamine uptake and [^{3}H]GBR 12783 binding to the dopamine uptake complex in the rat striatum, 335
Torda, T., see Saito, K., 35
Torrioli, G., see Rossini, P.M., 300
Traversa, R., see Rossini, P.M., 300
Tulipan, N.B., see Whetsell Jr., W.O., 18
Twyman, R.E., Rogers, C.J. and MacDonald, R.L., Pentobarbital and picrotoxin have reciprocal actions on single $GABA_A$ receptor channels, 89

Vanselow, J., see Bähr, M., 197
Villalba, R.M., see Martínez-Murillo, R., 121

Wakabayashi, K., Takahashi, H., Ohama, E. and Ikuta, F., Tyrosine hydroxylase-immunoreactive intrinsic neurons in the Auerbach's and Meissner's plexuses of humans, 259
Walsh, F.S., see Doherty, P., 1
Westlund, K.N., McNeill, D.L. and Coggeshall, R.E., Glutamate immunoreactivity in rat dorsal root axons, 13
Whetsell Jr., W.O., Allen, G.S. and Tulipan, N.B., Alteration of kainic acid and quinolinic acid toxicity by neostriatal transplants in vitro, 18
Wieloch, T., see Magnusson, K., 264
Woolf, N.J., Jacobs, R.W. and Butcher, L.L., The pontomesencephalotegmental cholinergic system does not degenerate in Alzheimer's disease, 277
Wroblewski, J.T., see Danysz, W., 340

Yamatsu, K., see Ikeda, M., 202
Yoshida, M., see Sakai, M., 157

Ziylan, Y.Z. and Ates, N., Age-related changes in regional patterns of blood–brain barrier breakdown during epileptiform seizures induced by pentylenetetrazol, 179